

## **Study Group Exclusion Criteria:**

This is a list of the conditions that might affect the FDP level in the blood and so the DR-70 test results. If these conditions are present, please use multiple consequent measurements of DR-70 level along with another marker applicable depending on the tumor type.

Note: A good portion of the conditions in the list is also affecting CEA test and that is why it is always good to use a panel of few tumor markers and multiple consequent measurements to be able to cover those conditions.

- Pregnant
- Undergoing pregnancy-related complications:
  - o Abruptio placenta
  - o Eclampsia
  - o Pre-eclampsia
  - o Intrauterine fetal death
- Inflammatory/autoimmune disease:
  - o Addison's
  - o Celiac disease
  - o Grave's disease
  - o Good pasture's syndrome
  - o HIV
  - o Hashimoto's Thyroiditis
  - o Lupus, Multiple Sclerosis
  - o Polymyalgia
  - o Raynaud's
  - o Scleroderma
  - o Sjogren's disease
- Surgery
- Trauma
- Burns
- Cirrhosis
- Renal disease/insufficiency
- Acute & Chronic Infections:
  - o Bacteremia
  - o Septicaemia (sepsis)
  - o Cellulitis
- Thrombotic problems:
  - o Pulmonary embolism (PE)
  - o Disseminated intravascular coagulation (DIC)
  - o Deep vein thrombosis (DVT)
  - o Myocardial infarction (MI)
  - o Ischemic heart disease
  - o Congenital heart disease
  - o Peripheral vascular disease
  - o Portacaval shunt
- Massive blood transfusion
- Therapy for coagulopathic conditions:
  - o Aspirin
  - o Ibuprofen (Nurofen, Advil, Mortin, Nuprin)
  - o Aggrenox
  - o Coumadin (Warfarin)
  - o Heparin
  - o Lovenox (Enoxaparin)
  - o Palvix
  - o Ibuprofen (Nurofen, Advil, Mortin, Nuprin)
- Hypercholesterolemia